

$$x = (R-r) \cos \frac{rt}{R} + c \cos \frac{(R-r)t}{R}, y = (R-r) \sin \frac{rt}{R} - c \sin \frac{(R-r)t}{R}, t \in R.$$

$$x = (R-r) \cos \varphi + c \cos \frac{(R-r)\varphi}{r}, y = (R-r) \sin \varphi - c \sin \frac{(R-r)\varphi}{r}, \varphi \in R.$$

$$x = (\sqrt{3}-1)r \cos \frac{t}{\sqrt{3}} + 2r \cos \frac{(\sqrt{3}-1)t}{\sqrt{3}}$$

$$y = (\sqrt{3}-1)r \sin \frac{t}{\sqrt{3}} - 2r \sin \frac{(\sqrt{3}-1)t}{\sqrt{3}}$$

$$t \in \langle 0; 10.7387\pi \rangle$$

$$R = \sqrt{3}r, c = 2r$$

$$x = (\sqrt{3}-1)r \cos \varphi + 2r \cos (\sqrt{3}-1)\varphi$$

$$y = (\sqrt{3}-1)r \sin \varphi - 2r \sin (\sqrt{3}-1)\varphi$$

$$\varphi \in \langle 0; 6.2\pi \rangle$$